LIST OF FIGURES

Figure 1.	Galveston Bay bait shrimping areas sampled by Bessette (1985)	14
Figure 2.	Mean salinities in Galveston Bay during 1984 sampling period (data from Bessette, 1985)	15
Figure 3.	Mean bycatch : shrimp ratio by area and month (data from Bessette, 1985)	16
Figure 4.	Monthly CPUE of dominant bycatch species in Trinity Bay (data from Bessette, 1985)	17
Figure 5.	Monthly CPUE of dominant bycatch species in upper Galveston Bay (data from Bessette, 1985)	18
Figure 6.	Monthly CPUE of dominant bycatch species in west Galveston Bay (data from Bessette, 1985)	19
Figure 7.	Monthly CPUE of dominant bycatch species in lower Galveston Bay (data from Bessette, 1985)	20
Figure 8.	Monthly CPUE of dominant bycatch species in East Bay (data from Bessette, 1985)	21
Figure 9.	Monthly CPUE of dominant bycatch species in Chocolate Bay (data from Bessette, 1985)	22
Figure 10.	Bycatch CPUE in Galveston Bay by area and month (data from Bessette, 1985)	23
Figure 11.	Bycatch as percent of total catch weight in Galveston Bay (data from Bessette, 1985)	24
Figure 12.	Shrimp CPUE in Galveston Bay by area and month (data from Bessette, 1985)	25
Figure 13.	Mean tow duration by area and month (data from Bessette, 1985)	26
Figure 14.	Percent of total bycatch (number and biomass) contributed by dominant fish and/or invertebrates during July and August 1981 (from Lamkin 1984)	27
Figure 15.	Percent of total bycatch (number and biomass) contributed by dominant fish and/or invertebrates during September and October 1981 (from Lamkin 1984)	28

Figure 16.	Percent of total bycatch (number and biomass) contributed by dominant fish and/or invertebrates during November and December 1981 (from Lamkin 1984)	29
Figure 17.	Percent of total bycatch (number and biomass) contributed by dominant fish and/or invertebrates during May and June 1982 (from Lamkin 1984)	30
Figure 18.	Average monthly brown and white shrimp catch from Galveston Bay during 1982-89 (data from NMFS landings files)	53
Figure 19.	Average total shrimp catch (with monthly %'s) from Galveston Bay during 1982-89 (data from NMFS landings files)	53
Figure 20.	Average monthly shrimp effort (days fished) for Galveston Bay during 1982-89 (with monthly %'s)	54
Figure 21.	Map of Galveston Bay showing designated fishing zones for sample collection	54
Figure 22.	Sample size estimates for three fishing areas in Galveston Bay	55
Figure 23.	Distribution of trawls sampled during 1992 data collection in Galveston Bay	56
Figure 24.	Monthly sampling intensity for the three "fishing zones" sampled in 1992	57
Figure 25.	Number of species collected (finfish and invertebrates) in the three fishing zones during new sampling efforts	57
Figure 26.	Size distribution (tail length) of white shrimp (<i>Penaeus setiferus</i>) in shrimp trawls sampled in A) Trinity Bay, B) Upper and East Galveston Bays and C) Lower and West Galveston bays	58
Figure 27.	Size distribution (tail length) of brown shrimp (<i>Penaeus aztecus</i>) in shrimp trawls sampled in A) Trinity Bay, B) Upper and East Galveston Bays and C) Lower and West Galveston bays	59
Figure 28.	Size distribution of gulf menhaden (<i>Brevoortia patronus</i>) in shrimp trawls sampled in A) Trinity Bay, B) Upper and East Galveston Bays and C) Lower and West Galveston bays	60
Figure 29.	Size distribution of Atlantic croaker (<i>Micropogonias undulatus</i>) in shrimp trawls sampled in A) Trinity Bay, B) Upper and East Galveston bays and C) Lower and West Galveston bays	61

Figure 30.	Size distribution of spot (<i>Leiostomus xanthurus</i>) in shrimp trawls sampled in A) Trinity Bay, B) Upper and East Galveston Bays and C) Lower and West Galveston bays	62
Figure 31.	Size distribution of cutlassfish (<i>Trichiurus lepturus</i>) in shrimp trawls sampled in A) Trinity Bay, B) Upper and East Galveston Bays and C) Lower and West Galveston bays	63
Figure 32.	Size distribution of sand seatrout (<i>Cynoscion arenarius</i>) in shrimp trawls sampled in A) Trinity Bay, B) Upper and East Galveston Bays and C) Lower and West Galveston bays	64
Figure 33.	Size distribution of spotted seatrout (<i>Cynoscion nebulosus</i>) in shrimp trawls sampled in A) Trinity Bay, B) Upper and East Galveston Bays and C) Lower and West Galveston bays	65
Figure 34.	Size distribution of red drum (<i>Sciaenops ocellatus</i>) in shrimp trawls sampled in A) Trinity Bay, B) Upper and East Galveston Bays and C) Lower and West Galveston bays	66
Figure 35.	Size distribution of southern flounder (<i>Paralichthys lethostigma</i>) in shrimp trawls sampled in A) Trinity Bay, B) Upper and East Galveston bays and C) Lower and West Galveston bays	67
Figure 36.	Size distribution of black drum (<i>Pogonias cromis</i>) in shrimp trawls sampled in A) Trinity Bay, B) Upper and East Galveston Bays and C) Lower and West Galveston bays	68
Figure 37.	Plot of finfish: shrimp ratios for the three fishing zones sampled during May - November 1992 (Lower Bay, Trinity Bay, Epper Bay)	31
Figure 38.	Plot of invertebrate: shrimp ratios for the three fishing zones sampled during May - November 1992 (Lower Bay, Trinity Bay, Epper Bay) 1	31

LIST OF TABLES

Table 1.	Temperature and salinity in Galveston Bay sampling areas during 1981-82 (from Lamkin 1984)	31
Table 2.	Biomass and percentage of total catch for bait shrimp and bycatch taken from Galveston Bay during July 1981 through June 1982 (from Lamkin 1984)	32
Table 3.	Monthly catch of southern flounder by commercial shrimp vessels in in various areas of Galveston Bay (data from Matlock, 1982)	33
Table 4.	Baywide frequency of fishes, May-November 1984 (data from Bessette, 1985)	34
Table 5.	Baywide bycatch species percentage by biomass and number (data from Bessette, 1985)	36
Table 6.	Occurrence of fish by species and bay area (data from Bessette 1985)	38
Table 7.	Baywide frequency of invertebrates, May-November 1984 (data from Bessette 1985)	45
Table 8.	Average monthly shrimp: fish ratios and associated 95% confidence intervals (± t.05 standard deviation) calculated from log (ln) transformed data (from Lamkin 1984)	46
Table 9.	Minimum sampling intensity projected for Galveston Bay bycatch study	69
Table 10.	Monthly distribution of samples taken in Galveston Bay 'fishing zones' during 1992.	69
Table 11.	Total CPUE (by number and weight) and temporal distribution of individual species captured in Galveston Bay shrimp trawls during 1992	70
Table 12.	Mean monthly catch per unit (CPUE) by number and biomass for all species captured in Galveston Bay during March-November 1992	75
Table 13.	Catch per unit effort (CPUE) statistics for all species captured during shrimp trawling within each area of Galveston Bay during March-November 1992	92
Table 14.	Dominant bycatch species (by number and weight) observed during 1992 sampling efforts in Galveston Bay	121

Table 15.	Listing of large, unusual or noteworthy items which were captured in trawls during 1992	122
Table 16.	Ratios of shrimp to finfish and invertebrates for numbers and weights of each species group captured in shrimp trawls by month and arean	
Table 17.	Estimated bycatch of finfish and invertebrates based on 1992 landings	126
Table 18.	Listing of individual species captured in fishery independent trawl surveys conducted by the Texas parks and Wildlife Department (79 species total)	132
Table 19.	Comparison and statistical analysis of length-frequency distributions for NMFS and TPWD data	134
Table 20.	Catch per unit effort for each species during 1992 NMFS sampling and TPWD fishery independent surveys	137
Table 21.	Ratio of finfish and invertebrate catch to shrimp landings from Texas Parks and Wildlife Department fishery independent surveys	145
Table 22.	Linear regression results for statistical (regression) comparison of NMFS and TPWD bycatch ratios	145
Appendix 1	Listing of common and scientific names of finfish and invertebrates captured during 1992 trawl bycatch study	169